

EFFECTIVE BASE POINT FREENESS ON A NORMAL SURFACE

TAKESHI KAWACHI

Department of Mathematics, Tokyo Institute of Technology

ABSTRACT. We prove effective base point freeness of the adjoint linear system on normal surfaces with a boundary.

0. INTRODUCTION

Let Y be a compact normal two dimensional projective variety over $\mathbb{C}$ (we call it “normal surface” for short). Let $y \in Y$ be a given point, and D be a nef and big $\mathbb{Q}$ -divisor on Y such that $K_Y + [D]$ is a Cartier divisor, There is various numerical criteria on D when the adjoint linear system $|K_Y + [D]|$ is free at y . The main theorem includes all earlier result and gives the first effective version if y is log-terminal singularity.

To describe the main theorem we prepare notation.

Definition. Let Y be a normal surface, and B be an effective $\mathbb{Q}$ -divisor on Y . Let y be a point on Y . Then the triple (Y, B, y) is said to be a germ of *quasi-log-terminal* singularity if the following conditions are satisfied:

- (1) $[B] = 0$, where $[B]$ is the integral part of B .
- (2) Let $f: X \rightarrow Y$ be the minimal resolution of y (if y is a smooth point then let f be the blowing-up at y). If we write

$$K_X = f^*(K_Y + B) + \sum a_j F_j \quad (\text{for } a_j \in \mathbb{Q})$$

then $a_j > -1$ whenever F_j is f -exceptional.

Remark. The differences with “log-terminal” are:

- (1) We don’t mention where $K_Y + B$ is $\mathbb{Q}$ -Cartier or not. The pull-back and the intersection are used by Mumford’s $\mathbb{Q}$ -valued pull-back and intersection theory.
- (2) f is only “minimal resolution”. Thus the $\mathbb{Q}$ -divisor $\sum a_j F_j$ may not be normal crossings. Hence “quasi-log-terminal” may not be “log-terminal” (cf. [KMM]).
- (3) If (Y, B, y) is quasi-log-terminal then $(Y, 0, y)$ which is ignored the boundary, is log-terminal.

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX

Let (Y, B, y) be a germ of normal surface singularity with a boundary B , and let $f: X \rightarrow Y$ be the minimal resolution of y (if y is a smooth point then the blowing-up at y). Let Z is the fundamental cycle of y (if y is a smooth point then the exceptional divisor of f). Let $\Delta_y = f^*K_Y - K_X$. Δ_y is an effective $\mathbb{Q}$ -divisor if y is singular, and $\Delta_y = -Z$ if y is smooth.

Definition.

$$\delta_y = \begin{cases} -(Z - \Delta_y)^2, & \text{if } y \text{ is quasi-log-terminal} \\ 0, & \text{if } y \text{ is not quasi-log-terminal} \end{cases}$$

Definition. We define

$$\mu(B, y) = \max\{\mu \mid \mu(Z - \Delta_y) \leq f^*B\}$$

Remark.

- (1) This δ_y satisfies $0 \leq \delta_y \leq 4$ ([KM]).
- (2) If y is a smooth point then $2\mu = \text{mult}_y B$. If y is quasi-log-terminal then $\mu < 1$.

The main result is the following.

Theorem. Let (Y, B) be a normal surface with a boundary B and assume that $[B] = 0$. Let M be a Cartier divisor on Y such that $M - (K_Y + B)$ is nef big. Let D be an effective $\mathbb{Q}$ -divisor on Y which is $\mathbb{Q}$ -linearly equivalent to $M - (K_Y + B)$. Let y be a point on Y . Let $\mu = \mu(B, y)$. Assume that $D^2 > (1 - \mu)^2 \delta_y$. If $DC \geq (1 - \mu)\delta_y/2$ for all curves C in Y such that $y \in C$ then the linear system $|M|$ is free at y . Furthermore if y is a singular point which is not of type A_n then $|M|$ is free at y .

We can get the following immediately.

Corollary. Let Y be a normal surface. Let D be a nef $\mathbb{Q}$ -divisor on Y such that $K_Y + [D]$ is Cartier. Let y be a point on Y . Let $B = [D] - D$ and $\mu = \mu(B, y)$. Assume that $D^2 > (1 - \mu)^2 \delta_y$. If $DC \geq (1 - \mu)\delta_y/2$ for all curves C in Y such that $y \in C$ then the linear system $|K_Y + [D]|$ is free at y .

Remark. If y is a smooth point or a rational double point then this theorem is essentially the theorem of Ein-Lazarsfeld ([EL]). If y is not a quasi-log-terminal point then the result are proved in [ELM]. If $B = 0$ then the result is described in [KM].

In section 1, we remark the facts about quasi-log-terminal singularity. These are used by the proof of the main theorem. In section 2, we prove the theorem.

1. FACTS ABOUT QUASI-LOG-TERMINAL SINGULARITY

Throughout this section, let (Y, B, y) be a germ of quasi-log-terminal surface singularity. Let $\Delta_y = f^*K_Y - K_X = \sum a_j \Delta_j$ where Δ_i is an irreducible exceptional divisor lying over y . Let $B = B_y + B'$ and $f^*B_y = \widetilde{B}_y + B_y^{\text{exc}} = \widetilde{B}_y + \sum b_j \Delta_j$ where B_y consists of the components which passing through y , B' is disjoint from y and $\widetilde{B}_y$ is the proper transformation of B_y . Let Z be the fundamental cycle of y . Hence we have

$$f^*(K_Y + B) = K_X + \widetilde{B}_y + \Delta_y + B_y^{\text{exc}} + f^*B'.$$

First we bound the value of $\mu(B, y)$. we recall the definition of $\mu(B, y)$.

$$\mu(B, y) = \max\{\mu \mid \mu(Z - \Delta_y) \leq f^*B\}$$

Lemma 1. *If (Y, B, y) is quasi-log-terminal then $\mu(B, y) < 1$.*

Proof. If $\mu(B, y) \geq 1$ then $B_y^{\text{exc}} \geq Z - \Delta_y$. Hence we have $\Delta_y + B_y^{\text{exc}} \geq Z$. On the other hand, since (Y, B, y) is quasi-log-terminal, all coefficients of $\Delta_y + B_y^{\text{exc}}$ must be less than 1. That is contradiction. $\square$

Since B is an effective divisor, B_y^{exc} is also effective. Therefore all coefficients of Δ_y are less than 1. This is the case of classical “log-terminal” surface singularity without boundary.

We defined $\delta_y = -(Z - \Delta_y)^2$. This is bounded by the following lemma (cf. [KM]).

Lemma 2. *$0 < \delta_y < 2$ if y is log-terminal but not a smooth or a rational double point.*

Proof. Since the intersection matrix $\|\Delta_i \Delta_j\|$ is negative definite and $Z - \Delta_y$ is effective and non-zero, we have $\delta_y > 0$. On the other hand, we have

$$\begin{aligned} (Z - \Delta_y)^2 &= Z(Z - \Delta_y) - \Delta_y(Z - \Delta_y) \\ &= Z(Z + K_X) + K_X(Z - \Delta_y) \\ &> Z(Z + K_X) = 2Pa(Z) - 2. \end{aligned}$$

Since the log-terminal singularity is the rational singularity, we have $Pa(Z) = 0$ by [A]. Therefore we have $\delta_y < 2$. $\square$

It is easy to prove that $\delta_y = 4$ if y is a smooth point and that $\delta_y = 2$ if y is a rational double point.

Let C be an irreducible divisor on Y and $y \in C$. Let $f^*C = \tilde{C} + \sum c_{i,j} \Delta_j$, and $\tilde{C} \Delta_i = 1$ for an i and $\tilde{C} \Delta_j = 0$ for $j \neq i$. That is, we assume that $\tilde{C}$ meets only Δ_i . Next, we study the value $a_i + c_{i,i}$. Since all log-terminal singularities are classified in [B], we examine it in each cases. we set indexes as in Appendix.

1.1 The case of type A_n . Let $A = A(w_1, \dots, w_n)$ be the intersection matrix of type A_n where $w_i = -\Delta_i^2$.

$$A(w_1, \dots, w_n) = \begin{pmatrix} -w_1 & 1 & 0 & 0 \\ 1 & -w_2 & \ddots & 0 \\ 0 & \ddots & \ddots & 1 \\ 0 & 0 & 1 & -w_n \end{pmatrix}.$$

Let $a(w_1, \dots, w_n) = \det A(w_1, \dots, w_n)$, and we define $a() = 1$ for convention.

Let $\tilde{A}_{ij}$ be the (i, j) -component of A^{-1} . Suppose $j < i$ then we have

$$\tilde{A}_{ij} = \frac{1}{\det A} (-1)^{i+j} \begin{vmatrix} -w_1 & 1 & & & & & & & 0 \\ & 1 & \ddots & & 1 & 0 & & & \\ 0 & & 1 & -w_{j-1} & 1 & 0 & 0 & & \\ & & 0 & 0 & 1 & -w_{j+1} & 1 & 0 & \\ & & & & 0 & \ddots & -w_{i-1} & 0 & \\ & & & & & 0 & 1 & 1 & 0 \\ & & & & & & 0 & -w_i & \ddots & 0 \\ & & & & & & 0 & 1 & \ddots & 1 \\ & & & & & & & 0 & 1 & -w_n \end{vmatrix}$$

$$= \frac{1}{\det A} (-1)^{i+j} a(w_1, \dots, w_{j-1}) a(w_{i+1}, \dots, w_n).$$

If $j > i$ then we have the following as same as above.

$$\tilde{A}_{ij} = \frac{1}{\det A} (-1)^{i+j} a(w_1, \dots, w_{i-1}) a(w_{j+1}, \dots, w_n).$$

Also we have

$$\tilde{A}_{ii} = \frac{1}{\det A} a(w_1, \dots, w_{i-1}) a(w_{i+1}, \dots, w_n).$$

Since

$$A(w_1, \dots, w_n) \begin{pmatrix} a_1 \\ \vdots \\ a_n \end{pmatrix} = \begin{pmatrix} 2 - w_1 \\ \vdots \\ 2 - w_n \end{pmatrix},$$

we have the following.

$$\begin{aligned} a_i &= \sum_{j=1}^{i-1} \tilde{A}_{ij} (2 - w_j) + \tilde{A}_{ii} (2 - w_i) + \sum_{j=i+1}^n \tilde{A}_{ij} (2 - w_j) \\ &= \frac{1}{\det A} \{ a(w_{i+1}, \dots, w_n) ((-1)^{i+1} (2 - w_1) + \dots + (-1)^{2i-1} a(w_1, \dots, w_{i-2}) (2 - w_{i-1})) + \\ &\quad a(w_1, \dots, w_{i-1}) a(w_{i+1}, \dots, w_n) (2 - w_i) + \\ &\quad a(w_1, \dots, w_{i-1}) ((-1)^{2i+1} a(w_{i+2}, \dots, w_n) (2 - w_{i+1}) + \dots + (-1)^{i+n} (2 - w_n)) \}. \end{aligned}$$

Since $a(w_1, \dots, w_j) = -w_j a(w_1, \dots, w_{j-1}) - a(w_1, \dots, w_{j-2})$, we have

$$\begin{aligned} &(-1)^{i+1} (2 - w_1) + (-1)^{i+2} a(w_1) (2 - w_2) + \dots + (-1)^{2i-1} a(w_1, \dots, w_{i-2}) (2 - w_{i-1}) \\ &= (-1)^{i+1} \{ 2 - w_1 + (-1) (2a(w_1) + a()) + a(w_1, w_2) \} + \dots + \\ &\quad (-1)^{i-2} (2a(w_1, \dots, w_{i-2}) + a(w_1, \dots, w_{i-3}) + a(w_1, \dots, w_{i-1})) \} \\ &= (-1)^{i+1} (1 + (-1)^{i-2} (a(w_1, \dots, w_{i-2}) + a(w_1, \dots, w_{i-1}))) \\ &= (-1)^{i+1} - a(w_1, \dots, w_{i-2}) - a(w_1, \dots, w_{i-1}). \end{aligned}$$

We also have the following by same calculation.

$$\begin{aligned} & (-1)^{2i+1}a(w_{i+2}, \dots, w_n)(2 - w_{i+1}) + \dots + (-1)^{i+n}(2 - w_n) \\ & = -a(w_{i+1}, \dots, w_n) - a(w_{i+2}, \dots, w_n) + (-1)^{i+n}. \end{aligned}$$

Hence we have

$$\begin{aligned} a_i &= \frac{1}{\det A} \{ a(w_{i+1}, \dots, w_n)((-1)^{i+1} - a(w_1, \dots, w_{i-2}) - a(w_1, \dots, w_{i-1})) + \\ & \quad a(w_1, \dots, w_{i-1})a(w_{i+1}, \dots, w_n)(2 - w_i) + \\ & \quad a(w_1, \dots, w_{i-1})((-1)^{i+n} - a(w_{i+1}, \dots, w_n) - a(w_{i+2}, \dots, w_n)) \} \\ &= \frac{1}{\det A} \{ (-1)^{i+1}a(w_{i+1}, \dots, w_n) + (-1)^{i+n}a(w_1, \dots, w_{i-1}) \\ & \quad - a(w_1, \dots, w_{i-2})a(w_{i+1}, \dots, w_n) - a(w_1, \dots, w_{i-1})a(w_{i+1}, \dots, w_n)w_i \\ & \quad - a(w_1, \dots, w_{i-1})a(w_{i+2}, \dots, w_n) \} \\ &= 1 + \frac{1}{\det A} \{ (-1)^{i+1}a(w_{i+1}, \dots, w_n) + (-1)^{i+n}a(w_1, \dots, w_{i-1}) \}. \end{aligned}$$

Since $(A(w_1, \dots, w_n)(c_{i,j}))_j = -1$ if $j = i$ and $= 0$ if $j \neq i$, we have

$$\begin{aligned} c_{i,j} &= -\tilde{A}_{ji} \\ &= \frac{1}{\det A} (-1)^{i+j+1} \begin{cases} a(w_1, \dots, w_{j-1})a(w_{i+1}, \dots, w_n), & \text{if } j \leq i \\ a(w_1, \dots, w_{i-1})a(w_{j+1}, \dots, w_n), & \text{if } j > i. \end{cases} \end{aligned}$$

Therefore we have the following.

$$\begin{aligned} a_i + c_{i,i} &= 1 + \frac{1}{|\det A|} (-|a(w_{i+1}, \dots, w_n)| - |a(w_1, \dots, w_{i-1})| + \\ & \quad |a(w_1, \dots, w_{i-1})||a(w_{i+1}, \dots, w_n)|) \\ &= 1 + \frac{1}{|\det A|} ((|a(w_1, \dots, w_{i-1})| - 1)(|a(w_{i+1}, \dots, w_n)| - 1) - 1) \end{aligned}$$

Hence we get the following proposition.

Proposition 1. *Let C as above. Assume y is of type A_n . If $n \geq 3$ and $i \neq 1, n$ then $a_i + c_{i,i} \geq 1$. If $i = 1$ or n then $a_i + c_{i,i} = 1 - 1/|\det A|$ (for $n \geq 1$).*

In the case of type A_n we have $\delta_y = 2 - a_1 - a_n$. Indeed, since $(\Delta_y - Z)\Delta_i = 2 - w_i - (-w_i + 2) = 0$ if $i \neq 1, n$ and $(\Delta_y - Z)\Delta_i = 1$ if $i = 1$ or n , we get the equation $\delta_y = 2 - a_1 - a_n$.

1.2 The case of type D_n . Let $D = D(w_1, \dots, w_{n-2})$ be the intersection matrix of type D_n where $w_i = -\Delta_i^2$.

$$D(w_1, \dots, w_n) = \begin{pmatrix} -w_1 & 1 & 0 & & & 0 \\ 1 & -w_2 & \ddots & & & \\ 0 & \ddots & \ddots & 1 & 0 & 0 \\ & 0 & 1 & -w_{n-2} & 1 & 1 \\ & & 0 & 1 & -2 & 0 \\ 0 & & 0 & 1 & 0 & -2 \end{pmatrix}.$$

Let $d(w_1, \dots, w_{n-2}) = \det D(w_1, \dots, w_{n-2})$, and we define $d() = 4$ for convention.

Let $\tilde{D}_{ij}$ be the (i, j) -component of D^{-1} . As same as of type A_n , we have

$$\tilde{D}_{ij} = \frac{1}{\det D} \begin{cases} (-1)^{i+j} a(w_1, \dots, w_{j-1}) d(w_{i+1}, \dots, w_{n-2}), & i \leq n-2, j \leq i \\ (-1)^{i+j} a(w_1, \dots, w_{i-1}) d(w_{j+1}, \dots, w_{n-2}), & i \leq n-2, i < j \leq n-2. \end{cases}$$

Let $i \leq n-2, j = n-1$. In this case we have

$$\begin{aligned} \tilde{D}_{i,n-1} &= \frac{1}{\det D} (-1)^{i+n-1} \begin{vmatrix} -w_1 & 1 & 0 & & & 0 \\ 1 & \ddots & 1 & 0 & & \\ 0 & \ddots & -w_{i-1} & 0 & & \\ & 0 & 1 & 1 & 0 & \\ & & 0 & -w_{i+1} & \ddots & 0 & 0 \\ & & 0 & 1 & -w_{n-2} & 1 & 1 \\ 0 & & & 0 & 1 & 0 & -2 \end{vmatrix} \\ &= \frac{1}{\det D} (-1)^{i+n-1} (-2) a(w_1, \dots, w_{i-1}). \end{aligned}$$

Also we have

$$\begin{aligned} \tilde{D}_{in} &= \tilde{D}_{n-1,i} = \tilde{D}_{ni} \\ &= \frac{1}{\det D} (-1)^{i+n-1} (-2) a(w_1, \dots, w_{i-1}). \end{aligned}$$

Now we suppose $i, j \geq n-1$. In this case, we have

$$\begin{aligned} \tilde{D}_{n-1,n-1} &= \tilde{D}_{n,n} \\ &= \frac{1}{\det D} \begin{vmatrix} -w_1 & 1 & 0 & 0 \\ 1 & \ddots & 1 & 0 \\ 0 & 1 & -w_{n-2} & 1 \\ 0 & 0 & 1 & -2 \end{vmatrix} = \frac{1}{\det D} a(w_1, \dots, w_{n-2}, 2). \end{aligned}$$

$$\begin{aligned}\tilde{D}_{n-1,n} &= \tilde{D}_{n,n-1} \\ &= \frac{1}{\det D}(-1) \begin{vmatrix} -w_1 & 1 & 0 & 0 \\ 1 & \ddots & 1 & 0 \\ 0 & 1 & -w_{n-2} & 1 \\ 0 & 0 & 1 & 0 \end{vmatrix} = \frac{1}{\det D} a(w_1, \dots, w_{n-3}).\end{aligned}$$

Note that there is an equation

$$d(w_1, \dots, w_{n-2}) = -4a(w_1, \dots, w_{n-2}, 1).$$

Now we calculate the value a_i . The calculation will proceed as same as of type A_n . But one thing differs, that is $d(w_{n-2}) = -4w_2 + 4$. For $i \leq n-2$, we have the following.

$$\begin{aligned}a_i &= \frac{1}{\det D} \{d(w_{i+1}, \dots, w_{n-2})((-1)^{i+1} - a(w_1, \dots, w_{i-2}) - a(w_1, \dots, w_{i-1})) + \\ &\quad a(w_1, \dots, w_{i-1})d(w_{i+1}, \dots, w_{n-2})(2 - w_i) \\ &\quad - a(w_1, \dots, w_{i-1})(d(w_{i+1}, \dots, w_{n-2}) + d(w_{i+2}, \dots, w_{n-2}))\} \\ &= \frac{1}{\det D} \{(-1)^{i+1}d(w_{i+1}, \dots, w_{n-2}) - a(w_1, \dots, w_{i-2})d(w_{i+1}, \dots, w_{n-2}) \\ &\quad - a(w_1, \dots, w_{i-1})d(w_{i+1}, \dots, w_{n-2})w_i - a(w_1, \dots, w_{i-1})d(w_{i+2}, \dots, w_{n-2})\} \\ &= 1 + \frac{1}{\det D}(-1)^{i+1}d(w_{i+1}, \dots, w_{n-2}) = 1 - \frac{1}{|\det D|}|d(w_{i+1}, \dots, w_{n-2})|.\end{aligned}$$

Since $c_{i,i} = -\tilde{D}_{ii}$, we have the following.

$$a_i + c_{i,i} = 1 + \frac{1}{|\det D|}|d(w_{i+1}, \dots, w_{n-2})|(|a(w_1, \dots, w_{i-1})| - 1).$$

For $i = n-1, n$, we have $a_{n-1} = a_n = a_{n-2}/2$. Hence we have

$$a_{n-1} = a_n = \frac{1}{2} + (-1)^{n-1} \frac{2}{\det D} = \frac{1}{2} - \frac{2}{|\det D|}.$$

Therefore we have the following.

$$\begin{aligned}a_{n-1} + c_{n-1,n-1} &= a_n + c_{n,n} \\ &= 1 + \frac{1}{\det D} \{2(-1)^{n-1} - \frac{1}{2}d(w_1, \dots, w_{n-2}) - a(w_1, \dots, w_{n-2}, 2)\} \\ &= 1 + \frac{1}{\det D} \{2(-1)^{n-1} + 2a(w_1, \dots, w_{n-2}, 1) - a(w_1, \dots, w_{n-2}, 2)\} \\ &= 1 + \frac{1}{\det D} \{2(-1)^{n-1} - 2a(w_1, \dots, w_{n-2}) - 2a(w_1, \dots, w_{n-3}) + \\ &\quad 2a(w_1, \dots, w_{n-2}) + a(w_1, \dots, w_{n-3})\} \\ &= 1 + \frac{1}{\det D} \{2(-1)^{n-1} - a(w_1, \dots, w_{n-3})\} \\ &= 1 + \frac{1}{|\det D|}(|a(w_1, \dots, w_{n-3})| - 2) \geq 1.\end{aligned}$$

Hence we get the following proposition.

Proposition 2. *Let C as above. If y is of type D_n then $a_i + c_{i,i} \geq 1$ for all i .*

1.3 The case of type E_n . If the dual graph is in the Appendix then we write $(m; a, b, c; d, e)$ or may write $(m; a, b, c; d, e; -2)$. We set indexes i as in the Appendix. There are only 15 types in all by the classification ([B]). Hence we can calculate $a_i + c_{i,i}$ directly. The result is in Appendix.

Hence we get the following proposition.

Proposition 3. *Let C as above. If y is of type E_n then $a_i + c_i > 1$ for all i .*

2. PROOF OF THE THEOREM

We recall notation. Let (Y, B) be a normal surface with a boundary B such that $[B] = 0$. Let M be a Cartier divisor on Y . Let D be an effective $\mathbb{Q}$ -divisor on Y , $\mathbb{Q}$ -linearly equivalent to the nef big divisor $M - (K_Y + B)$. Let y be a point of Y . Let $f: X \rightarrow Y$ be the global minimal resolution, if y is a smooth point, f factors the blowing-up at y . Let $\Delta = f^*K_Y - K_X$. Let Δ_y be the components of Δ supported on $f^{-1}(y)$, and $\Delta' = \Delta - \Delta_y$. Let Z be the fundamental cycle of y . Let $\mu = \mu(B, y)$, where $\mu(B, y) = \max\{\mu \mid \mu(Z - \Delta_y) \leq f^*B\}$. Let $\delta_y = -(Z - \Delta_y)^2$ if y is quasi-log-terminal, and $\delta_y = 0$ if y is not quasi-log-terminal. Let $B = B_y + B'$ where B_y consists of the component which contain y , and B' is disjoint from y . Let $f^*B_y = \widetilde{B}_y + B_y^{\text{exc}}$ where $\widetilde{B}$ is the proper transformation of B_y , and $f(B_y^{\text{exc}}) = \{y\}$. Since $M \sim K_Y + D + B$ is Cartier, $f^*(K_Y + D + B) = K_X + f^*(D + B) + \Delta$ is integral. Hence $D + B$ is also integral coefficients, thus $\lceil D \rceil = D + B$.

First we treat the case which y is not quasi-log-terminal. In this case, we assume that $D^2 > 0$ and $DC \geq 0$ for all curve $C \ni y$, because we defined $\delta_y = 0$. Since D is nef and big, this always hold. Hence it is sufficient to prove that y is not a base point of $|M|$.

Since f^*D is nef and big, we have $H^1(X, K_X + \lceil f^*D \rceil) = 0$ by Sakai's lemma [S1] where $\lceil \rceil$ means the ‘‘round-up’’. On the other hand,

$$\begin{aligned} K_X + \lceil f^*D \rceil &= \lceil K_X + f^*D \rceil = \lceil f^*(K_Y + B) - \Delta - f^*B + f^*D \rceil \\ &= \lceil f^*(K_Y + \lceil D \rceil) - (\Delta_y + B_y^{\text{exc}}) - \Delta' - f^*B' \rceil \\ &= f^*(K_Y + \lceil D \rceil) - \lfloor \Delta_y + B_y^{\text{exc}} \rfloor - \lfloor \Delta' + f^*B' \rfloor. \end{aligned}$$

Since y is not quasi-log-terminal, the round-down $\lfloor \Delta_y + B_y^{\text{exc}} \rfloor \neq 0$. Let $S = \lfloor \Delta_y + B_y^{\text{exc}} \rfloor$ and $T = \lfloor \Delta' + f^*B' \rfloor$. Hence we have a surjection

$$H^0(X, f^*(K_Y + \lceil D \rceil) - T) \rightarrow H^0(S, (f^*(K_Y + \lceil D \rceil) - T)|_S) \rightarrow 0.$$

Since $f(S) = \{y\}$, $\mathcal{O}_S((f^*(K_Y + \lceil D \rceil) - T)|_S)$ is trivial. Hence we get a section of $\mathcal{O}_X(f^*(K_Y + \lceil D \rceil) - T)$ which does not vanish at any point of $\text{Supp } S$. Therefore we get a global section of $\mathcal{O}_X(f^*(K_Y + \lceil D \rceil))$ which does not vanish on $f^{-1}(y)$ by multiplying the global section of $\mathcal{O}_X(T)$.

From now we assume that y is quasi-log-terminal. Let $D = D_y + D'$ where all irreducible divisors in D_y contain y and D' is the others. Let $f^*D_y = \widetilde{D}_y + D_y^{\text{exc}} = \sum d_i D_i + \sum d'_j \Delta_j$.

Let $f^*B_y = \sum b_i D_i + \sum b'_j \Delta_j$. Let $\Delta_y = \sum a_j \Delta_j$. Since $D^2 > -((1 - \mu)(Z - \Delta_y))^2$, $f^*D - (1 - \mu)(Z - \Delta_y)$ is big. Hence if necessary, we change D by $\mathbb{Q}$ -linearly equivalence and we assume $D_y \neq 0$. We define the number

$$c = \min \left\{ \frac{1 - b_i}{d_i}, \frac{1 - a_j - b'_j}{d'_j} \right\}$$

Since $d_i, d'_j > 0$, $0 \leq b_i < 1$ and $a_j + b'_j < 1$ we have $c > 0$.

Lemma 3. $c \leq 1/2$.

Proof. Since $D^2 > -((1 + \varepsilon)(1 - \mu)(Z - \Delta_y))^2$ for sufficiently small ε , we have $D - (1 + \varepsilon)(1 - \mu)(Z - \Delta_y) > 0$. Hence we have $d'_j - (1 - \mu)(z_j - a_j) > 0$ where $Z = \sum z_j \Delta_j$. Thus $d'_j + a_j - z_j + \mu(z_j - a_j) > 0$. By the definition of μ , we have $b'_j \geq \mu(z_j - a_j)$. Hence we have $d'_j + a_j - z_j + b'_j > 0$. Therefore $d'_j + a_j + b'_j > z_j \geq 1$ because z_j is positive integer. On the other hand, Since $K_Y + [D] = K_Y + D + B$ is Cartier, $f^*D + f^*B + \Delta = f^*(K_Y + D + B) - K_X$ is integral divisor. Hence $d'_j + a_j + b'_j$ is also integral. Since $d'_j + a_j + b'_j \geq 2$, we have $2(1 - a_j/2 - b'_j/2) \leq d'_j$. Thus we have

$$c \leq \frac{1 - a_j - b'_j}{d'_j} \leq \frac{1}{2} - \frac{a_j + b'_j}{2d'_j} \leq \frac{1}{2}. \quad \square$$

Let

$$\begin{aligned} R &= f^*D - cf^*D = f^*[D] + \Delta - f^*B - cf^*D - \Delta \\ &= f^*[D] + \Delta - \left(\sum (b_i + cd_i)D_i + \sum (b'_j + cd'_j + a_j)\Delta_j \right) - (f^*B' + f^*D' + \Delta'). \end{aligned}$$

Since $f^*[D] + \Delta = f^*(K_Y + [D]) - K_X$ is Cartier,

$$\begin{aligned} [R] &= f^*[D] + \Delta - \left[\sum (b_i + cd_i)D_i + \sum (b'_j + cd'_j + a_j)\Delta_j \right] - \lfloor f^*B' + f^*D' + \Delta' \rfloor \\ &= f^*[D] + \Delta - (D_1 + \cdots + D_s) - E' - Q \end{aligned}$$

where $\lfloor \sum (b_i + cd_i)D_i \rfloor = D_1 + \cdots + D_s$, $\lfloor \sum (b'_j + cd'_j + a_j)\Delta_j \rfloor = E'$, and $\lfloor f^*B' + f^*D' + \Delta' \rfloor = Q$.

Suppose $E' \neq 0$. Since $R = (1 - c)f^*D$ is nef big and $D_1, \dots, D_s$ are not appear in the fractional part of R , we have $H^1(X, K_X + [R] + D_1 + \cdots + D_s) = 0$ by Kawamata-Viehweg vanishing theorem. Hence we have the surjection

$$H^0(X, f^*(K_Y + [D]) - Q) \rightarrow H^0(E', (f^*(K_Y + [D]) - Q)|_{E'}) \rightarrow 0.$$

Since $f(E') = \{y\}$ and Q is disjoint from y , we get a global section of $K_Y + [D]$ which is not vanish at y .

Now we assume $E' = 0$. Note that the minimality of c induce that $b_i + cd_i \leq 1$. Thus the minimal value c is obtained by $(1 - b_i)/d_i$, hence $s \geq 1$. By Kawamata-Viehweg vanishing theorem, we have

$$H^0(X, K_X + \lceil R \rceil + D_2 + \cdots + D_s) = H^0(K_X + f^* \lceil D \rceil + \Delta - D_1 - Q) = 0.$$

Thus there is the surjection

$$H^0(X, K_X + f^* \lceil D \rceil + \Delta - Q) \rightarrow H^0(D_1, K_{D_1} + (f^* \lceil D \rceil + \Delta - D_1 - Q)|_{D_1}) \rightarrow 0.$$

So it is sufficient to prove $\lceil R \rceil D_1 > 1$. Indeed, if $\lceil R \rceil D_1 > 1$ then $\lceil R \rceil D_1 \geq 2$ because $\lceil R \rceil$ is Cartier. Thus $(f^* \lceil D \rceil + \Delta - D_1 - Q)D_1 \geq \lceil R \rceil D_1 \geq 2$. Hence we get the nowhere vanishing global section in $H^0(D_1, K_{D_1} + (f^* \lceil D \rceil + \Delta - D_1 - Q)|_{D_1})$. Hence we get the global section of $\mathcal{O}(f^*(K_Y + \lceil D \rceil))$ which is not vanish on D_1 , especially not vanish at y .

Hence we prove $\lceil R \rceil D_1 > 1$. Remark that

$$\lceil R \rceil = R + \left\{ \sum_{i \neq 1, \dots, s} (b_i + cd_i)D_i + \sum (b'_j + cd'_j + a_j)\Delta_j + (\text{others}) \right\}$$

where $\{(\text{divisor})\}$ is the fractional part of (divisor). Since $E' = 0$, $\{\sum (b'_j + cd'_j + a_j)\Delta_j\} = \sum (b'_j + cd'_j + a_j)\Delta_j$. Therefore

$$\lceil R \rceil D_1 \geq (1 - c)f^*DD_1 + \sum (b'_j + cd'_j + a_j)\Delta_j D_1.$$

Thus it is sufficient to prove

$$(1) \quad (1 - c)(1 - \mu)\delta_y/2 + \sum (b'_j + cd'_j + a_j)\Delta_j D_1 > 1.$$

If D_1 meets more components of Δ_y then $\sum (b'_j + cd'_j + a_j)\Delta_j D_1$ is larger. Thus we assume $\Delta_i D_1 = 1$ for an i and $\Delta_i D_1 = 0$ for $j \neq i$. Let $f^*f_*D_1 = D_1 + \sum c_{i,j}\Delta_j$. Since $f^*B_y + cf^*D_y = D_1 + (\text{others})$, we have $c_{i,j} \leq b'_j + cd'_j$. But by Proposition 1 through 3, $c_{i,i} + a_i \geq 1$ if y is of type D_n or of type E_n , and if y is of type A_n and $i \neq 1, n$. They contradict to the assumption $E' = 0$. Thus we assume y is of type A_n and $j = 1, n$ or y is a smooth point.

Thus if y is of type D_n or E_n and $D^2 > (1 - \mu)^2\delta_y$ then y is not a base point of $|M|$.

First we assume that y is a smooth point or of type A_1 . In this case, $b'_1 = \mu(1 - a_1)$. Let $p = \mu(D, y)$, hence $d'_1 = p(1 - a_1)$. Let $w_1 = -\Delta_1^2$. Then we have $a_1 = 1 - 2/w_1$ and $\delta_y = 4/w_1$. Hence $\delta_y = 2(1 - a_1)$. Since $f^*D - (1 - \mu)(Z - \Delta_y)$ is big, we have $p + \mu > 1$ as in the proof of lemma 3. Hence we have

$$\begin{aligned} \lceil R \rceil D_1 &\geq (1 - c)(1 - \mu)\frac{\delta_y}{2} + (b'_1 + cd'_1 + a_1) \\ &= (1 - \mu)\frac{\delta_y}{2} + b'_1 + a_1 + (d'_1 - (1 - \mu)\frac{\delta_y}{2})c \\ &= (1 - \mu)(1 - a_1) + \mu(1 - a_1) + a_1 + (p(1 - a_1) - (1 - \mu)(1 - a_1))c \\ &= 1 + (p + \mu - 1)(1 - a_1)c > 1. \end{aligned}$$

Next we assume that y is of type A_n and $n \geq 2$. We may change the indexes of Δ_j , we assume $a_1 \leq a_n$. We also assume that D_1 meets Δ_i where $i = 1$ or n .

Case 1: We assume that $i = n$ or $a_1 = a_n$.

It is sufficient to prove $(1 - c)(1 - \mu)\delta_y/2 + b'_i + cd'_i + a_i > 1$ by the inequality of (1). Since the definition of μ , we have $\mu(1 - a_i) \leq b'_i$. Since $f^*D - (1 - \mu)(Z - \Delta_y)$ is big, we have $d'_i > (1 - \mu)(1 - a_i)$. Since $\delta_y = 2 - a_1 - a_n$, we have

$$\begin{aligned}
& (1 - c)(1 - \mu)\delta_y/2 + b'_i + cd'_i + a_i \\
& > (1 - \mu) \left(1 - \frac{a_1 + a_n}{2} \right) + \mu(1 - a_i) + a_i + \\
& \quad \left((1 - \mu)(1 - a_i) - (1 - \mu) \left(1 - \frac{a_1 + a_n}{2} \right) \right) c \\
& = 1 - \frac{a_1 + a_n}{2} + a_i + \mu \left(1 - a_i - 1 + \frac{a_1 + a_n}{2} \right) + c(1 - \mu) \left(1 - a_i - 1 + \frac{a_1 + a_n}{2} \right) \\
& = 1 + \frac{a_n - a_1}{2} - \mu \frac{a_n - a_1}{2} - c(1 - \mu) \frac{a_n - a_1}{2} \quad (\text{let } i = n) \\
& = 1 + (1 - c)(1 - \mu) \frac{a_n - a_1}{2} \geq 1.
\end{aligned}$$

Case 2: We assume that $i = 1$ and $a_1 < a_n$.

Let $B_y = bD_1 + B''$ and $\text{ord}_{\Delta_i} f^*B'' = b''_i$. Since

$$\sum (b'_i + cd'_i + a_i)\Delta_i = (f^*B_y + cf^*D_y + \Delta_y)^{\text{exc}} \geq (f^*f_*D_1 + f^*B'' + \Delta_y)^{\text{exc}},$$

we have $b'_i + cd'_i + a_i \geq a_i + c_i + b''_i$. Since $a_1 + c_{1,1} = 1 - 1/|\det A|$, if $b''_1 \geq 1/|\det A|$ then $[R]D_1 > 1$ because $\delta_y > 0$.

Suppose that $b''_1 < 1/|\det A|$. Let $B'' = \sum b_k D_k$ and $f^*D_k = \widetilde{D}_k + \sum (c_{k_1,j} + \cdots + c_{k_{r_k},j})\Delta_j$ if $\widetilde{D}_k$ meets $\Delta_{k_1}, \dots, \Delta_{k_{r_k}}$ counted with intersection multiplicity. Since

$$\frac{1}{|\det A|} > b''_1 = \sum_{k,i} b_k c_{k_i,1} \geq \sum_k b_k c_{n,1} r_k = \frac{\sum b_k r_k}{|\det A|},$$

we have $\beta = \sum b_k r_k < 1$. By the definition of μ we have

$$\begin{aligned}
\mu & \leq \frac{b''_n}{1 - a_n} = \frac{bc_{1,n} + \sum b_k c_{k_i,n}}{1 - a_n} \leq \frac{bc_{1,n} + \sum b_k r_k c_{n,n}}{1 - a_n} \\
& = \frac{bc_{1,n} + \beta c_{n,n}}{1 - a_n}.
\end{aligned}$$

Hence we have

$$\begin{aligned}
& (1-c)(1-\mu)\frac{\delta_y}{2} \\
& \geq (1-c) \left(1 - \frac{bc_{1,n} + \beta c_{n,n}}{1-a_n}\right) \frac{\delta_y}{2} \\
& = (1-c) \frac{\delta_y}{2(1-a_n)} ((1-a_n)|\det A| - b - \beta a(w_1, \dots, w_{n-1})) \frac{1}{|\det A|} \\
& = (1-c) \frac{\delta_y}{2(1-a_n)} ((1-b) + (1-\beta)a(w_1, \dots, w_{n-1})) \frac{1}{|\det A|}.
\end{aligned}$$

If $b \leq \beta$ then

$$\begin{aligned}
(1-c)(1-\mu)\frac{\delta_y}{2} & \geq (1-c) \frac{\delta_y}{2(1-a_n)} (1-\beta) \frac{1 + a(w_1, \dots, w_{n-1})}{|\det A|} \\
& = (1-c)(1-\beta) \frac{\delta_y}{2(1-a_n)} (1-a_n) \\
& = (1-c)(1-\beta) \frac{\delta_y}{2}.
\end{aligned}$$

Since $c \leq 1/2$ by lemma 3, we have

$$\begin{aligned}
& (1-c)(1-\mu)\frac{\delta_y}{2} + \frac{\beta}{|\det A|} \\
& \geq (1-\beta) \frac{\delta_y}{4} + \frac{\beta}{|\det A|} \\
& = \frac{1}{|\det A|} \left(\frac{1}{4} (2 + a(w_1, \dots, w_{n-1}) + a(w_2, \dots, w_n)) (1-\beta) + \beta \right) \\
& \geq \frac{1}{|\det A|} \left(\frac{3}{2} - \frac{1}{2}\beta \right) \\
& > \frac{1}{|\det A|}.
\end{aligned}$$

Hence we have

$$\begin{aligned}
& (1-c)(1-\mu)\frac{\delta_y}{2} + b'_1 + cd_1 + a_1 \\
& \geq (1-c)(1-\mu)\frac{\delta_y}{2} + a_1 + c_{1,1} + \sum b_k c_{k_i,1} \\
& \geq (1-c)(1-\mu)\frac{\delta_y}{2} + 1 - \frac{1}{|\det A|} + \frac{\beta}{|\det A|} \\
& > 1.
\end{aligned}$$

Finally we assume that $\beta < b$. In this case we have

$$\begin{aligned}
(1-c)(1-\mu)\frac{\delta_y}{2} &\geq \frac{\delta_y}{4(1-a_n)}((1-b) + (1-\beta)a(w_1, \dots, w_{n-1}))\frac{1}{|\det A|} \\
&= \frac{\delta_y}{4(1-a_n)}(1-\beta)(1+a(w_1, \dots, w_{n-1}))\frac{1}{|\det A|} - \frac{\delta_y}{4(1-a_n)}\frac{1}{|\det A|}(b-\beta) \\
&> (1-\beta)\frac{\delta_y}{4} - \frac{\delta_y}{4}\frac{1}{1+a(w_1, \dots, w_{n-1})}(1-\beta) \\
&= (1-\beta)\frac{\delta_y}{4}\left(1 - \frac{1}{1+a(w_1, \dots, w_{n-1})}\right) \\
&\geq (1-\beta)\frac{\delta_y}{6}.
\end{aligned}$$

Hence we have

$$\begin{aligned}
(1-c)(1-\mu)\frac{\delta_y}{2} + \frac{\beta}{|\det A|} &> (1-\beta)\frac{\delta_y}{6} + \frac{\beta}{|\det A|} \\
&= \frac{1}{|\det A|}\left(\frac{1}{6}(2+a(w_1, \dots, w_{n-1})+a(w_2, \dots, w_n))(1-\beta)+\beta\right) \\
&\geq \frac{1}{|\det A|}.
\end{aligned}$$

Therefore we have

$$(1-c)(1-\mu)\frac{\delta_y}{2} + b'_1 + cd_1 + a_1 > 1.$$

Thus we have $[R] D_1 > 1$, we complete the proof. $\square$

APPENDIX

type	dual graph	$a_i + c_i$	$x =$
1	$(m; 2, 2; 2, 2)$	$\left(1 + \frac{5}{x}, 1 + \frac{1}{3x}, \frac{4}{3} + \frac{2}{x}, \frac{4}{3} + \frac{2}{x}, 1 + \frac{1}{3x}, 1 + \frac{1}{x}\right)$	$6m - 11$
2	$(m; 2, 2; 3)$	$\left(1 + \frac{5}{3x}, 1 + \frac{1}{9x}, \frac{4}{3} + \frac{2}{3x}, 1 + \frac{1}{9x}, 1 + \frac{1}{3x}\right)$	$2m - 3$
3	$(m; 3; 3)$	$\left(1 + \frac{5}{x}, 1 + \frac{1}{3x}, 1 + \frac{1}{3x}, 1 + \frac{1}{x}\right)$	$6m - 7$
4	$(m; 2, 2, 2; 2, 2)$	$\left(1 + \frac{11}{x}, 1 + \frac{1}{2x}, \frac{3}{2} + \frac{5}{2x}, \frac{3}{2} + \frac{6}{x}, \frac{4}{3} + \frac{14}{3x}, 1 + \frac{1}{x}, 1 + \frac{5}{2x}\right)$	$12m - 23$
5	$(m; 2, 2, 2; 3)$	$\left(1 + \frac{11}{x}, 1 + \frac{1}{2x}, \frac{3}{2} + \frac{5}{2x}, \frac{3}{2} + \frac{6}{x}, 1 + \frac{1}{x}, 1 + \frac{5}{2x}\right)$	$12m - 19$
6	$(m; 4; 2, 2)$	$\left(1 + \frac{11}{x}, 1 + \frac{1}{2x}, \frac{4}{3} + \frac{1}{4}3x, 1 + \frac{1}{x}, 1 + \frac{5}{2x}\right)$	$12m - 17$
7	$(m; 4; 3)$	$\left(1 + \frac{11}{x}, 1 + \frac{1}{2x}, 1 + \frac{1}{x}, 1 + \frac{5}{2x}\right)$	$12m - 13$
8	$(m; 2, 2, 2, 2; 2, 2)$	$\left(1 + \frac{29}{x}, 1 + \frac{1}{x}, \frac{8}{5} + \frac{22}{5x}, \frac{9}{5} + \frac{51}{5x}, \frac{8}{5} + \frac{92}{5x}, \frac{4}{3} + \frac{38}{3x}, 1 + \frac{3}{x}, 1 + \frac{7}{x}\right)$	$30m - 59$
9	$(m; 2, 2, 2, 2; 3)$	$\left(1 + \frac{29}{x}, 1 + \frac{1}{x}, \frac{8}{5} + \frac{22}{5x}, \frac{9}{5} + \frac{51}{5x}, \frac{8}{5} + \frac{92}{5x}, 1 + \frac{3}{x}, 1 + \frac{7}{x}\right)$	$30m - 49$
10	$(m; 2, 3; 2, 2)$	$\left(1 + \frac{29}{x}, 1 + \frac{1}{x}, \frac{6}{5} + \frac{22}{5x}, \frac{4}{3} + \frac{38}{3x}, 1 + \frac{3}{x}, 1 + \frac{7}{x}\right)$	$30m - 47$
11	$(m; 2, 3; 3)$	$\left(1 + \frac{29}{x}, 1 + \frac{1}{x}, \frac{6}{5} + \frac{22}{5x}, 1 + \frac{3}{x}, 1 + \frac{7}{x}\right)$	$30m - 37$
12	$(m; 3, 2; 2, 2)$	$\left(1 + \frac{29}{x}, 1 + \frac{1}{x}, \frac{7}{5} + \frac{51}{5x}, \frac{4}{3} + \frac{38}{3x}, 1 + \frac{3}{x}, 1 + \frac{7}{x}\right)$	$30m - 53$
13	$(m; 3, 2; 3)$	$\left(1 + \frac{29}{x}, 1 + \frac{1}{x}, \frac{7}{5} + \frac{51}{5x}, 1 + \frac{3}{x}, 1 + \frac{7}{x}\right)$	$30m - 43$
14	$(m; 5; 2, 2)$	$\left(1 + \frac{29}{x}, 1 + \frac{1}{x}, \frac{4}{3} + \frac{38}{3x}, 1 + \frac{3}{x}, 1 + \frac{7}{x}\right)$	$30m - 41$
15	$(m; 5; 3)$	$\left(1 + \frac{29}{x}, 1 + \frac{1}{x}, 1 + \frac{3}{x}, 1 + \frac{7}{x}\right)$	$30m - 31$

REFERENCES

- [A] M. Artin, *On isolated rational singularities of surfaces*, Amer. J. Math. **88** (1966), 129–136.
- [B] E. Brieskorn, *Rationale singularitäten komplexer flächen*, Invent. Math. **14** (1968), 336–358.
- [EL] L. Ein–R. Lazarsfeld, *Global generation of pluricanonical and adjoint linear series on projective threefolds*, Jour. AMS **6** (1993), 875–903.
- [ELM] L. Ein, R. Lazarsfeld, V. Mařek, *Global generation of linear series on terminal threefolds*, International Journal of Math. **6** (1995), 1–18.
- [K] T. Kawachi, *On freeness theorem of the adjoint bundle on a normal surface*, alg-geom/9603022.
- [KM] T. Kawachi, V. Mařek, *Reider–Type Theorems on Normal Surfaces*, Journal of Algebraic Geometry (submitting).
- [KMM] Y. Kawamata–K. Matsuda–K. Matsuki, *Introduction to the Minimal Model Problem*, Adv. Studies in Pure Math. **10** (1987), 283–360.
- [R] I. Reider, *Vector bundles of rank 2 and linear systems on algebraic surfaces*, Ann. Math. **127** (1988), 309–316.
- [S1] F. Sakai, *Weil divisors on normal surfaces*, Duke Math. J. **51** (1984), 877–887.
- [S2] ———, *Reider–Serrano’s method on normal surfaces*, Algebraic Geometry, Proceedings: L’Aquila 1988, Lect. Notes in Math. **1417** (1990), 301–319.

2-12-1 OH-OKAYAMA, MEGURO-KU, TOKYO, JAPAN, 152
E-mail address: kawachi@math.titech.ac.jp